

Simulating Quantum Field Theories on a Quantum Computer

Stephen Jordan



JOINT CENTER FOR
QUANTUM INFORMATION
AND COMPUTER SCIENCE



UNIVERSITY OF
MARYLAND

Can quantum computers simulate all physical processes efficiently?

Universality Conjecture:

Quantum circuits can simulate all physical dynamics in $\text{poly}(E, V, t, 1/\epsilon)$ time.


Status:

Non-relativistic QM	Yes: Now being optimized
Quantum Field Theories	Probably: In progress
Quantum Gravity/Strings	Nobody knows


Quantum Field Theory

- Much is known about using quantum computers to simulate quantum systems.
- Why might quantum field theory be different?
 - Field has infinitely many degrees of freedom
 - Relativistic
 - Particle number not conserved
 - Formalism looks different.


When do we need QFT?




Nuclear Physics



Accelerator Experiments




Cosmic Rays



Coarse-grained many-body systems


2015 NERSC Usage By Discipline

(MPP HOURS IN MILLIONS)




2015 NERSC Usage By Discipline

(MPP HOURS IN MILLIONS)




Classical Algorithms

Feynman diagrams



Break down at strong coupling or high precision

Lattice methods




Cannot calculate scattering amplitudes

There's room for exponential speedup by quantum computing.


A QFT Computational Problem

Input: a list of momenta of incoming particles.


Output: a list of momenta of outgoing particles.



S-matrix



Particle
accelerator



Results So Far


- Efficient quantum simulation algorithms:

	Bosonic	Fermionic
Massive	Jordan, Lee, Preskill <i>Science</i> , 336:1130 (2012)	Jordan, Lee, Preskill <i>ArXiv:1404.7115</i> (2014)
Massless	?	?

- BQP-hardness: classical computers cannot perform certain QFT simulations efficiently
[S. Jordan, H. Krovi, K. Lee, K. Preskill, 2017]
- Better Speed and Broken Symmetries
[A. Moosavian and S. Jordan, 2017]

Representing Quantum Fields

A field is a list of values, one for each location in space.



A quantum field is a superposition over classical fields.

$$\frac{1}{\sqrt{2}} \left| \begin{array}{c} \text{[Classical Field 1]} \end{array} \right\rangle - \frac{i}{\sqrt{2}} \left| \begin{array}{c} \text{[Classical Field 2]} \end{array} \right\rangle$$

A superposition over bit strings is a state of a quantum computer.

ϕ^4 -theory

Lagrangian density

$$\mathcal{L} = \frac{1}{2} \partial^\mu \phi \partial_\mu \phi - m^2 \phi^2 - \frac{\lambda}{4!} \phi^4$$

For quantum simulation we prefer
Hamiltonian formulation (equivalent)

$$H = \int d^d x [\pi^2 + (\nabla \phi)^2 + m^2 \phi^2 + \lambda \phi^4]$$

$$[\phi(x), \pi(y)] = i \delta^{(d)}(x - y)$$

Our Algorithms

1) Choose a lattice discretization.

Bound discretization error (by renormalization group).

2) Prepare physically realistic initial state.

Is the most time-consuming step.

This depends strongly on which QFT simulated.

3) Implement time-evolution by a quantum circuit.



Use Trotter formulae.

4) Perform measurements on final state.


Complicated by vacuum entanglement.

Lattice Cutoff

Continuum QFT = limit of a sequence of theories on successively finer lattices.



... continuum




Mass: m

Interaction strength: λ



Coarse grain




Mass: m'


Interaction strength: λ'

Lattice Cutoff

Continuum QFT = limit of a sequence of theories on successively finer lattices.




m and λ are functions of lattice spacing!



$$H = \frac{1}{2} \sum_{x \in \Omega} a^d [\pi^2 + (\nabla \phi)^2 + m^2 \phi^2 + \lambda \phi^4]$$



Coarse grain




$$H_{\text{eff}} = \frac{1}{2} \sum_{x \in \Omega'} (2a)^d [\pi^2 + (\nabla' \phi)^2 + m_{\text{eff}}^2 \phi^2 + \lambda_{\text{eff}} \phi^4 + g \phi^6 + \dots]$$

Simulation converges as a^2

Adiabatic State Preparation

$$H(s) = H_{\text{free}} + sH_{\text{interaction}}$$


Prepare wavepackets in free theory, then adiabatically turn on interaction. **Problem:**



Adiabatic State Preparation


Solution: intersperse backward time evolutions with time-independent Hamiltonians.

This winds back dynamical phase on each eigenstate without undoing adiabatic change of basis.



Simulating Detectors

- Measure energy in localized regions:



- Need smooth envelope function to avoid excessive vacuum noise!

$$H_f = \sum_{\mathbf{x}} f(\mathbf{x}) \mathcal{H}(\mathbf{x})$$


Runtimes

Weak Coupling:

$d = 1$	$(1/\epsilon)^{1.5}$
$d = 2$	$(1/\epsilon)^{2.376}$
$d = 3$	$(1/\epsilon)^{5.5}$

Strong Coupling:

	$\lambda_c - \lambda_0$	p	n_{out}
$d = 1$	$\left(\frac{1}{\lambda_c - \lambda_0}\right)^9$	p^4	n_{out}^5
$d = 2$	$\left(\frac{1}{\lambda_c - \lambda_0}\right)^{6.3}$	p^6	$n_{\text{out}}^{7.128}$



Fermions:

- Fermion doubling problem
- Free vacuum different from Bosonic case


Gross-Neveu:

$$H = \int dx \left[\sum_{j=1}^N \bar{\psi}_j \left(m_0 - i\gamma^1 \frac{d}{dx} \right) \psi_j + \frac{g^2}{2} \left(\sum_{j=1}^N \bar{\psi}_j \psi_j \right)^2 \right]$$

Fermion Doubling Problem


$$\frac{d\psi}{dx} \rightarrow \frac{\psi(x+a) - \psi(x-a)}{2a}$$

$$\sqrt{p^2 + m^2} \rightarrow \sqrt{\sin^2 p + m^2}$$



Wilson Term

$$H \rightarrow H - \frac{r}{2a} \sum_x \bar{\psi} (\psi(x+a) - 2\psi(x) + \psi(x-a))$$



Improved State Prep: Bosons

- In some cases (e.g. weakly coupled $d=2$), preparing the free vacuum is the rate limiting step.
- We can do this much faster using Bogoliubov transformation that looks like a Fast Fourier Transform.
[Somma, Jordan, unpublished]
- Essentially same idea as 2nd quantized FFT from:
[Babbush, Wiebe, McClean, McLain, Neven, Chan, 2017]

Improved State Prep: Fermions

- Two problems with adiabatic state preparation:
 - Cannot reach symmetry-broken phase
 - Runtime bound not practical: $O(\epsilon^{-8})$
- A solution for both:
 - First, prepare the vacuum from MPS
 - Then, resonantly excite single-particle wavepackets
 - Tighter analysis: CFT entropy and Floquet theory:
 $O(\epsilon^{-3.23})$

[A. Moosavian, S. Jordan, 2017]

Tensor Network Ansatzes





image credit: G. Evenbly

Tensor Network Ansatzes



Near-Term Prospects?

quantum
supremacy

science
applications

commercial
applications



- Simulating conformal field theories using MERA-based variational eigensolvers
- Simulating commuting Hamiltonians
- Simulating high-connectivity systems, e.g. spin glasses or SYK model

Near-Term Prospects?

quantum
supremacy

science
applications

commercial
applications



- Simulating conformal field theories using MERA-based variational eigensolvers
- Simulating commuting Hamiltonians
- Simulating high-connectivity systems, e.g. spin glasses or SYK model

Thanks!